

Photocatalytic reduction of CO₂ to C₁ fuels by (Ni/ZnO)@C nanoreactors

1st M.-L. Liu

Department of Environmental
Engineering
National Cheng Kung University
Tainan, Taiwan
morning.liu@aiasec.net

2nd L.-W. Wei

Department of Environmental
Engineering
National Cheng Kung University
Tainan, Taiwan
wei3532607@gmail.com

3rd H.-P. Wang

Department of Environmental
Engineering
National Cheng Kung University
Tainan, Taiwan
wanghp@ncku.edu.tw

Abstract

The (Ni/ZnO)@C core-shell nanoparticles were prepared by carbonization of Ni²⁺- and Zn²⁺-cyclodextrin complexes at 723 K for 2 h. ZnO and Ni encapsulated in carbon-shell were etched partially to form the (Ni/ZnO)@C yolk-shell nanoreactors for photocatalytic reduction of CO₂ to C₁ fuels. By XRD, it is clear that ZnO is the main zinc crystallite in the nanoreactors, and its nanoparticle size is between 10-20 nm. The TEM images of the nanoreactors indicate that Ni and ZnO having the nanosizes of 5-30 nm are capsulated in the porous carbon-shell that allows molecules to diffuse in and out for photocatalytic reduction of CO₂ to C₁ fuels. It is worth noting that ZnO in the (Ni/ZnO)@C yolk-shell nanoreactor plays the main photoactive role in photocatalytic degradation of methylene blue. However, excess Ni encapsulated in carbon-shell leads to a de-activity in photocatalytic degradation of MB and reduction of CO₂. By *in situ* FTIR spectroscopy, the disappearance of CO₂ is at the expense of formation of species containing CH and carbonyl groups, possibly related to yields of C₁ species such as HCOOH.

Keywords: CO₂ reduction, photocatalysis, nanoreactors, yolk-shell, methanol.