

Photocatalytic reduction of CO₂ to methanol by Cu₂O/TiO₂ heterojunctions

1st S.-P. Cheng

Department of Environmental
Engineering
National Cheng Kung University
Tainan, Taiwan
pingc2102@gmail.com

2nd L.-W. Wei

Department of Environmental
Engineering
National Cheng Kung University
Tainan, Taiwan
wei3532607@gmail.com

3rd H.-P. Wang

Department of Environmental
Engineering
National Cheng Kung University
Tainan, Taiwan
wanghp@ncku.edu.tw

Abstract

Conversion of CO₂ to low-carbon fuels using solar energy is considered an economically attractive and environmental friendly route. The development of novel catalysts and the use of solar energy via photocatalysis is the key to achieve the goal of chemically reducing CO₂ under mild conditions. Thus, in this study, the novel Cu₂O/TiO₂ heterojunctions were used for CO₂-to-low-carbon fuels. The p-n heterojunction is able to enhance the separation of photogenerated electron-hole pairs. By UV-vis diffuse reflection absorption spectroscopy, it is clear that Cu₂O coupled with TiO₂ causes a red-shift to the visible light range. Under a 6-h UV-vis irradiation, 12.4-70.6 μmol methanol/g-catalyst can be generated by the Cu₂O/TiO₂ heterojunctions. However, excess Cu₂O in the Cu₂O/TiO₂ heterojunctions may cause less absorption of UV-vis light and decrease the excited electrons from TiO₂, which may obstruct the photoactivity for reduction of CO₂.

Keywords: photocatalysis, CO₂, Cu₂O, TiO₂,
p-n heterojunction, methanol.